La ricerca su base di popolazione in Ticino

A.Bordoni
Registro Tumori Canton Ticino
Istituto Cantonale di Patologia, Locarno

Fondazione ticinese ricerca cancro
IOSI, Bellinzona
21.03.2012

Incidenza in Ticino e zone limitrofe

Confronti di incidenza dei tumori della prostata. Tassi standardizzati sulla popolazione Mondiale (per 100.000 abitanti)
Fonte: Cancer incidence in Five Continents, volume IX (1998-2002)

Regione	Tasso
Ferrara	41.5
Reggio Emilia	42.7
Firenze	46.1
Parma	49.1
Ticino	49.6
Torino	56.1
Milano	56.2
Biella	57.7
Veneto	57.9
Varese	63.0
Sondrio	67.6
Grigioni e Glarona	67.6
Vallese	75.1
San Gallo - Appenzello	76.9
Neuchatel	77.7
Svizzera	81.2
Ginevra	85.4
Vaud	88.7

I dati presentati indicano che l'incidenza nel nostro Cantone è relativamente bassa rispetto a quanto osservato nel resto della Svizzera e tende ad assomigliare ai dati osservati nel nord Italia

Asbesto e Mesotelioma

- Difficile da curare
- Diagnosi precoce ha poco senso
- Malattia debellabile mediante la prevenzione primaria, unica misura sicura

Mesotelioma

Incidenza in Ticino

Figura 1. Distribuzione delle diagnosi di mesotelioma secondo l'anno di incidenza e il sesso. Ticino, 1996-2007.

Mesotelioma

Incidenza Ticino a confronto

Figura 2. Confronti internazionali dei tassi di incidenza standardizzati per età sulla popolazione mondiale (per 100,000), uomini. Fonte: Cancer Incidence in Five Continents, vol IX.

Incidenza in Ticino e zone limitrofe

Incidenza nel mondo L'Agencia Internazionale di Ricerca sul Cancro (IARC) di Lione (WHO),

Tassi di incidenza standardizzati sulla popolazione mondiale (per 100.000)
TUMORI DELLA PROSTATA - Uomini

Trend di incidenza in Ticino e Svizzera, Prostata

Trend di incidenza per classi di età Ticino, 1996-2007, Prostata

Numero di pazienti e età media Ticino, 1996-2008, Prostata

Età media alla diagnosi (IC 95%)

Chernobyl

Chernobyl

European age-standardized incidence rates (EASR) for thyroid cancer observed by Swiss cancer registries during the period 1996–1998. GE, Geneva; NE, Neuchâtel; VD, Vaud; GA, St. Gall-Appenzell; BA, Basel; GG, Graubünden-Glarus; VS, Valais; TI, Ticino. Incidence of Total/papillary/other thyroid cancer by region.

Chernobyl

Observed and fitted European age-standardized incidence rates (EASR, per 100,000, age 0-85+) of thyroid cancer in pooled Swiss cancer registries (1980-1999), linear regression modelling

Chernobyl

Unexpected additional increase in the incidence of thyroid cancer among a recent birth cohort in Switzerland

Fabio Montanaro^a, Pierre Pury^b, Andrea Bordoni^a, Jean-Michel Lutz^{b,c}
and the Swiss Cancer Registries Network*

Data collected by the Swiss Cancer Registries Network (ASRT/VSKR) have been used to analyse trends in thyroid cancer during the last available 20 years, to make within-country geographical comparisons for current incidence rates. Age-standardized (European population) incidence rates per 100,000 for all morphologies combined ranges from 1.62 to 2.99 among males and from 2.13 to 8.09 among females in Switzerland. Regression analyses for both sexes combined detected an increase in time for papillary cases and a decrease for other types. Age-period-cohort analyses revealed that the youngest cohorts of men and women born after 1940 had an increased risk of all types of thyroid cancer while the cohort of people born between 1920 and 1939 were at increased risk of the papillary subtype. Assuming a higher sensitivity to ionizing radiation among the youngest people, a Chernobyl effect cannot be

definitively excluded and continuous study of this topic should be encouraged. *European Journal of Cancer Prevention* 15:178–186 © 2006 Lippincott Williams & Wilkins.

European Journal of Cancer Prevention 2006, 15:178–186

Keywords: thyroid cancer, incidence, time trends, Chernobyl accident, radiations, cancer registration

^aTicino Cancer Registry, Institute of Pathology, Locarno, ^bAssociation of Swiss Cancer Registries and ^cGeneva Cancer Registry, Switzerland

Correspondence to Dr Jean-Michel Lutz, Association Suisse des Registres des Tumeurs, Registre genevois des tumeurs, 55 Bd de la Cluse, CH-1205 Geneva, Switzerland
Tel: +41 22 379 49 50; fax: +41 22 379 49 71;
e-mail: jean-michel.lutz@imsp.unige.ch

Received 4 March 2005 Accepted 4 July 2005

Chernobyl o altro?

INCREASING TREND OF THYROID CANCER: A POPULATION-BASED STUDY

TUMOUR SUBTYPE	NO	I.C. (%)	NO	I.C. (%)	NO	I.C. (%)
anaplastic carcinoma	8	3.4%	4	7.5%	4	2.2%
other	10	4.2%	1	1.9%	9	4.9%

Figure 3 – Incidence trend of thyroid cancer by age group. Ticino, 1996-2007, females.

Age group	APC
20-49	13.1 * (5.1; 21.6)
50-69	0.3 (-6.6; 7.8)
>69	-0.7 (-8.9; 5.8)

Figure 4 – Incidence trend of papillary thyroid cancer by tumour size group. Ticino, 1996-2007, males and females.

Size group	APC
1-10 mm	8.9 * (0.3; 18.3)
11-20 mm	7.1 (-3.4; 18.8)
21-40 mm	13.1 (-8.6; 40.0)
>40 mm	0.9 (-37.9; 84.0)

Sex	APC (95% CI)
Men	4.4 (-22.2; 40.1)
ALL	6.7* (1.8; 11.8)

Figure 5 – Cause-specific survival of papillary thyroid cancer by size group. Ticino, 1996-2007, males and females.

Size group	5-year survival	10-year survival
1-10 mm	100%	100%
11-20 mm	96.7%	96.7%
21-40 mm	92%	92%
>40 mm	77.1%	61.7%

BAC

The inc change small s than a Epiden inciden diamet

PAT

All thyr were c accordi Change to dete

CONCLUSIONS

The widespread use of ultrasound and image-guided fine needle aspiration, and changes in tumour classification seem to be recognised factors of the increase of thyroid cancer incidence, particularly for small size tumours and papillary subtype. However, as reported recently in the literature, we detected a significant incidence increase in women, particularly in younger age and across all tumour size groups, suggesting that a real incidence increase could be present. Other factors (e.g. environmental agents and molecular pathways) could be involved in the observed pattern, so that additional European population-based data are needed in order to confirm these results.

Tumori polmonari

Inquinamento atmosferico, radon, tabagismo

Trend di incidenza dei tumori polmonari in Canton Ticino.
Tassi standardizzati per età sulla popolazione mondiale (per 100'000 abitanti).
Uomini, donne, uomini e donne, 1996-2010.

Tumori polmonari

Trend di incidenza dei tumori polmonari in Canton Ticino, per distretto.
Tassi standardizzati per età sulla popolazione mondiale (per 100'000 abitanti).
Uomini, 1996-2010.

DISTRETTO di DOMICILIO	APC annual change percentage
LOCARNO	-3.1 * (IC95%: -5.5; -0.6)
BELLINZONA	-2.1 (IC95%: -4.8; 0.8)
LUGANO	-1.5 (IC95%: -3.7; 0.8)
MENDRISIO	-3.8 * (IC95%: -6.2; -1.3)
TICINO	-2.2 * (IC95%: -3.5; -0.9)

* statisticamente significativo

Tumori polmonari

Trend di incidenza dei tumori polmonari in Canton Ticino, per distretto.
Tassi standardizzati per età sulla popolazione mondiale (per 100'000 abitanti).
Donne, 1996-2010.

DISTRETTO di DOMICILIO	APC annual percentage change
LOCARNO	4.5 * (IC95%: 0.8; 8.4)
BELLINZONA	4.5 * (IC95%: 0.4; 8.7)
LUGANO	1.6 (IC95%: -2.4; 5.8)
MENDRISIO	6.9 * (IC95%: 1.6; 12.6)
TICINO	3.6 * (IC95%: 1.1; 6.2)

* statisticamente significativo

* statisticamente significativo

Hindawi Publishing Corporation
 Journal of Cancer Epidemiology
 Volume 2011, Article ID 275758, 7 pages
 doi:10.1155/2011/275758

Research Article

Impact of Histopathological Diagnosis with Ancillary Immunohistochemical Studies on Lung Cancer Subtypes Incidence and Survival: A Population-Based Study

Bordoni Andrea,¹ Bongiovanni Massimo,² Mazzucchelli Luca,² and Spitale Alessandra¹

¹ Ticino Cancer Registry, Institute of Pathology, 6600 Locarno, Switzerland

² Division of Clinical Pathology, Institute of Pathology, 6600 Locarno, Switzerland

Correspondence should be addressed to Bordoni Andrea, andrea.bordoni@ti.ch

Received 26 September 2011; Accepted 7 November 2011

Academic Editor: Carmen J. Marsit

Copyright © 2011 Bordoni Andrea et al. This is an open access article distributed under the Creative Commons Attribution License, which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Purpose. The aim of this study was to assess the impact of immunohistochemical- (IHC-) studies on incidence and survival of lung cancer histotypes. *Patients and Methods.* Lung cancers occurred in southern Switzerland between 1996 and 2010 were selected by the Ticino Cancer Registry and categorised into adenocarcinoma (AC), squamous-cell-carcinoma (SqCC), small-cell-carcinoma (SmCC), and large-cell carcinoma/non-small-cell lung cancer (LCC/NSCLC). Incidence rates, annual-percentage-change (APC), and two-year overall survival (OS) (follow-up: 31.12.2010) were performed. *Results.* 2467 cases were selected: 997 (40.4%) AC; 522 (21.2%) LCC/NSCLC, 378 (15.3%) SmCC, and 570 (23.1%) SqCC. Trend-analysis showed significant increase in AC (APC: 4.6; 95% CI: 3.1; 6.0) and decrease of LCC/NSCLC, with significant joinpoint in 2003 (APC: -14.7; 95% CI: -21.6; -7.1). Improved OS and decreased OS were detected in SqCC and LCC/NSCLC, respectively. *Conclusions.* This study highlights that diagnosis with ancillary immunohistochemical studies will change incidence and survival of precisely defined lung cancer subtypes. It calls attention to the need for cautious interpretation of studies and clinical trials, where the diagnosis was based on histology unaccompanied by IHC studies, and to the need of standardised diagnostic procedures.

tasso standardizzato per età (Mondo)

age change

6.2; 14.8)

0.9; 9.6)

-5.9; 4.9)

-26.6; -3.3)

* statisticamente significativo

Come sono raccolti i dati in un Registro Tumori e con quali standard?

- Un Registro Tumori è un servizio pubblico adibito alla raccolta e archiviazione, analisi e interpretazione dei dati biologici tumorali dei pazienti
- Un Registro Tumori di popolazione raccoglie informazioni complete su tutti i nuovi casi che insorgono in una popolazione definita (es. Ticino)
- Garantire qualità e confrontabilità nel tempo e comparabilità internazionale

Come sono raccolti i dati in un Registro Tumori e con quali standard?

Come sono raccolti i dati in un Registro Tumori e con quali standard?

Come sono raccolti i dati in un registro tumori e con quali standard?

Come sono raccolti i dati in un registro tumori e con quali standard?

Es. del Ticino

WHO histological classification of tumours of the liver and intrahepatic bile ducts

Epithelial tumours		Miscellaneous Tumours	
Benign		Solitary fibrous tumour	8815/0
Hepatocellular adenoma (liver cell adenoma)	8170/0 ¹	Teratoma	9080/1
Focal nodular hyperplasia		Yolk sac tumour (endodermal sinus tumour)	9071/3
Intrahepatic bile duct adenoma	8160/0	Carcinosarcoma	8980/3
Intrahepatic bile duct cystadenoma	8161/0	Kaposi sarcoma	9140/3
Biliary papillomatosis	8264/0	Rhabdoid tumour	8963/3
		Others	
Malignant		Haemopoietic and lymphoid tumours	
Hepatocellular carcinoma (liver cell carcinoma)	8170/3	Secondary tumours	
Intrahepatic cholangiocarcinoma (peripheral bile duct carcinoma)	8160/3	Epithelial abnormalities	
Bile duct cystadenocarcinoma	8161/3	Liver cell dysplasia (liver cell change)	
Combined hepatocellular and cholangiocarcinoma	8180/3	Large cell type (large cell change)	
Hepatoblastoma	8970/3	Small cell type (small cell change)	
Undifferentiated carcinoma	8020/3	Dysplastic nodules (adenomatous hyperplasia)	
		Low-grade	
Non-epithelial tumours		High-grade (atypical adenomatous hyperplasia)	
Benign		Bile duct abnormalities	
Angiomyolipoma	8860/0	Hyperplasia (bile duct epithelium and peribiliary glands)	
Lymphangioma and lymphangiomatosis	9170/0	Dysplasia (bile duct epithelium and peribiliary glands)	
Haemangioma	9120/0	Intraepithelial carcinoma (carcinoma in situ) 8500/211	
Infantile haemangioendothelioma	9130/0		
Malignant		Miscellaneous lesions	
Epithelioid haemangioendothelioma	9133/1	Mesenchymal hamartoma	
Angiosarcoma	9120/3	Nodular transformation (nodular regenerative hyperplasia)	
Embryonal sarcoma (undifferentiated sarcoma)	8991/3	Inflammatory pseudotumour	
Rhabdomyosarcoma	8900/3		
Others			

¹Morphology code of the International Classification of Diseases for Oncology (ICD-O) (542) and the Systematized Nomenclature of Medicine (<http://snomed.org>). Behaviour is coded /0 for benign tumours, /1 for unspecified, borderline or uncertain behaviour, /2 for in situ carcinomas and grade III intraepithelial neoplasia and /3 for malignant tumours.

Come sono raccolti i dati in un registro tumori e con quali standard?

Es. del Ticino

Come sono raccolti i dati in un registro tumori e con quali standard?

Es. del Ticino

The screenshot shows a FileMaker database form for cancer registration. The form is organized into several sections:

- Identification and Registration:** Fields for patient ID, tumor RDT range, registration date, and global code.
- Personal Data:** Fields for name, address, sex, date of birth, and nationality.
- Diagnosis and Staging:** Fields for primary site, morphology, and ICD-10 codes for primary, terminal, and concomitant diseases.
- Treatment and Outcome:** Fields for date of death, nature of death, date of last modification, and survival status.
- Genetic and Molecular Data:** Fields for gene amplification (FISH), receptor status (estrogen, progesterone), and other molecular markers.

Perché servono i dati nominativi del paziente?

38'000 i pazienti contenuti nel database
2'400 ca i nuovi casi censiti ogni anno
40 le variabili considerate per ogni paziente
100'000 le variabili immesse ogni anno
70'000 i dossier medici considerati

10 il numero medio che un dossier viene aggiornato

Patologia

IOSI

Cliniche (Moncucco, St. Anna)

Oncologi sul territorio

La vicinanza alla patologia ci ha permesso di sviluppare sinergie

Histol Histopathol

<http://www.histo>

original article

Annals of Oncology 20: 628–635, 2009
doi:10.1093/annonc/mdn675
Published online 12 December 2008

Epidemiol
gastrointest
based s

Breast cancer classification according to immunohistochemical markers: clinicopathologic features and short-term survival analysis in a population-based study from the South of Switzerland

Paola Mazzola
¹Ticino Cancer Registry
Locarno, Switzerland

A. Spitale^{1*}, P. Mazzola¹, D. Soldini², L. Mazzucchelli² & A. Bordoni¹

¹Ticino Cancer Registry and ²Clinical Pathology, Institute of Pathology, Locarno, Switzerland

Received 27 August 2008; accepted 15 September 2008

Background: Breast cancer may be classified into distinct molecular subtypes based on gene expression profiling and/or immunophenotypic characteristics. Aim of the study was to investigate prevalence, clinicopathologic features and overall survival (OS) of molecular subtypes, in a large European population-based study.

Patients and methods: All invasive breast cancers from 2003 to 2007 were selected from the files of Ticino Cancer Registry. Molecular subtypes were defined by immunohistochemical markers. Clinicopathological characteristics and short-term OS were analyzed.

Results: Of 1214 invasive breast cancers, 73.2% were luminal A subtype, 13.8% luminal B, 7.4% basal like and 5.6% Her2/neu. Basal like presented largely in premenopausal women and displayed aggressive features, such as large tumor size, poorly differentiated cancers, high Ki-67 proliferation index and the worst 24-month OS. Luminal A included the highest percentage of patients >70, the highest proportion of stage I tumors and well/moderately differentiated lesions. Her2/neu was more frequent in postmenopausal women and showed the highest percentage of positive lymph nodes and stage IV cases.

Conclusion: This is a comprehensive European population-based study on breast cancer molecular subtypes. We provide strong evidence that the molecular classification is useful for clinical management and superior to World Health Organization classification in terms of short-term prognostic value.

Key words: cancer registry, immunohistochemistry, molecular breast cancer subtypes

Summary. Intraepithelial
(GISTs) are characterized
or *PDGFRA* mutations
raised major interest in
studies are still
diagnosed in 50% of
identified using
c-kit and PDGFR
features were

La vicinanza alla clinica ci ha permesso di sviluppare sinergie

Europa

Impatto
adv
stud
Aless
Anton

Preoper
widely ac
cancer (I
populati
overall s
LARC's d
and 2007
Ticino Ca

according to the first administered treatment: preoperative

Tumori, 95: 303-310, 2009

Prognosis of acute myeloid leukemia in the general population of southern Switzerland

Erika Lerch¹, Vittoria Espinasse¹, Giancarlo Scali², Oreste M. Geronzi¹ and Michele Ghielmini¹

¹Istituto Oncologico della Svizzera Italiana, ²Registrazione dei Tumori del Canton Ticino

ABSTRACT

Aims and background. To evaluate the prognostic factors, time to progression and overall survival in acute myeloid leukemia in the Italian population of southern Switzerland.

Methods and study design. A retrospective analysis of 100 acute myeloid leukemia patients from January 1984 to December 2007. Univariate and multivariate analysis for prognostic factors were performed.

Results. The incidence of acute myeloid leukemia in southern Switzerland is 2.6/100,000 per year. The median age at diagnosis was 66 years (range, 18 to 94). The median overall survival was 6 months, with a 2-year overall survival rate of 10%.

chapter 1

Annals of Oncology 18 (Supplement 1): i3-i6, 2007
doi:10.1093/annonc/mdl443

chapter 1

Epidemiology of hematological malignancies

D. Rodriguez-Abreu¹, A. Bordoni² & E. Zucca¹

¹IOSI, Oncology Institute of Southern Switzerland, Medical Oncology Department, Ospedale San Giovanni, Bellinzona; ²Registro cantonale dei tumori del Canton Ticino, Istituto cantonale di patologia, Locarno, Switzerland

incidence rates

Hematological malignancies comprise a collection of heterogeneous conditions, all originating from cells of the bone marrow and the lymphatic system. There are three major groups: leukemia, lymphoma, and plasma cell neoplasms. In general, in Western countries the overall incidence of hematological malignancies appears to be rising but it is very difficult to describe in a consistent and uniform way their epidemiological behavior, especially in Europe. The number of European Union (EU) Member States has increased and this expansion enlarged the Union to incorporate a diversity of peoples with a much larger degree of heterogeneity in lifestyle habits and disease risk, as well as in cancer incidence and mortality (either overall or site-specific). Moreover, there is not a unique European reporting system or registry for hematological cancers, so the estimation of the exact number of patients is difficult. In addition, the comparison of the incidences reported by various European registries may be biased because their populations can be predominantly either urban or rural, or because there are differences in the methodology and accuracy of registration.

The number of new patients in Europe diagnosed with hematological malignancies in 2005 can be estimated in ~230 000 patients. These new cases of leukemia, Hodgkin's and non-Hodgkin's lymphoma (NHL) and myeloma account for ~8% of all the new cancer patients diagnosed in Europe and the estimated deaths from these tumors account

for 5 and 6%; 21% between 65 and 74; 23% between 75 and 84; and 10% after 85 years of age.

Leukemias are usually divided into four major categories, with different clinical features and prognosis: acute lymphocytic leukemia, chronic lymphocytic leukemia, acute myelogenous leukemia, and chronic myelogenous leukemia.

The incidence rates for all types of leukemia are slightly higher among males than among females. There are also geographical and ethnic variations in leukemia rates. In the United States, the incidence is higher in Caucasians than in Afro-Americans and Hispanics. The American Indians/Alaskan natives have the lowest incidence rates. Trends in overall incidence of leukemia have generally been stable or slowly increasing. A substantial reduction in death rates from acute lymphoblastic leukemia (ALL), particularly in childhood, however, have been observed since the 1970s, thanks to advances in treatment and subsequent improvement in survival.

acute myelogenous leukemia (AML)

AML is mainly an adult's disease with a median age at presentation of 64 years. It accounts for ~30% of all leukemias in adults, and ~18 000 new patients are diagnosed in Europe each year, representing ~0.6% of all cancers. The annual incidence rate in Europe ranges from two per 100 000/year to four per 100 000/year. In the past decade, the trend in overall incidence of AML has generally been stable or slowly increasing in most European countries. Incidence in England and Wales,

Contributi tribuna medica

SEZIONE SCIENTIFICA - JOURNAL CLUB

SEZIONE SCIENTIFICA

SEZIONE SCIENTIFICA

SEZIONE SCIENTIFICA

SEZIONE SCIENTIFICA

Il Journal Club
to curato dal
Canton Ticino

**SOPRAVVIVUTI
TUMORE IN
CONTINENTALI
UNO STUI
DI POPOLAZIONE
LIVELLO N
(CONCORRERE)**

Coleman MP et al.
Tumors: a worldwide
(CONCORD). *Lancet*
9:730-756

Riassunto/Adatta
Dr.med. Andrea Bc
Dr.stat. Alessandra
Registro Tumori Cc
Istituto Cantonale

**LA DIAGNOSI DEL
CARCINOMA DELLA
PROSTATA
È REALMENTE IN
AUMENTO IN TICINO?**

A. Bordini, A. Spitale

Il carcinoma della prostata è uno
dei tumori più frequenti del maschio.
so viene considerato come un tumore
essenzialmente imprevedibile,
quanto può sviluppare metastasi il
pendentemente dall'accrescimento
del tumore, come pure mostrare il
gressione simultanea del tumore
della prostata e della vescicola

**INDICATORI
CLINICO-EPIDEMIOLOGICI
PER LA VALUTAZIONE
DELLA STRATEGIA
DI SCREENING
MAMMOGRAFICO
OPPORTUNITÀ
IN CANTONE**

A. Bordini, L. Mazzucchi

Introduzione
Il dibattito sui
mammografi

**INDICATORI
NEL TRATTAMENTO
DEL CARCINOMA
COLORETTO
IMPORTANZA
RACCOLTA
NUMERO
LINFONODI**

A. Spitale, L. Mazzucchi

Introduzione
Il carcinoma
colorettale è una
neoplasia delle
vie digerenti e
rappresenta la
seconda causa di
morte per cancro
in Ticino. Ogni
anno in Ticino
si registrano 70
di

**NUOVE ACQUISIZIONI
NEL TRATTAMENTO
DELLE NEOPLASIE
INTRACRANICHE:
GLI ASTROCITOMI.
COSA RESTA DA FARE?**

G. Pesce, D. Boscherini, V. Martin, A. Salatino,
S. Mauri, A. Spitale, A. Bordini, A. Richetti,
S. Tschuor, R. Wytttenbach, L. Mazzucchi,
R. Renella

Riassunto
I tumori primitivi del sistema nervoso
centrale (SNC) sono relativamente rari,

e di garantire ai pazienti migliori risultati
terapeutici.

Come si vedrà, per garantire un adeguato
approccio nel processo decisionale
diagnostico e terapeutico, in neuro-oncologia
più che in altre branche della medicina,
risulta determinante il coordinamento
degli sforzi di diversi servizi specialistici.

Le tecnologie e gli specialisti operanti
in Ticino, coadiuvati dalle relazioni con
altri centri di ricerca a livello nazionale
ed internazionale, permettono ai pazienti
di questo Cantone l'accesso ai migliori
standard terapeutici oggi esistenti.

1. Introduzione

Negli ultimi dieci anni l'interesse per la
neuro-oncologia è andato crescendo,
in parte perché le neoplasie cerebrali
più frequenti, vale a dire i gliomi

glioblastomi, sono state individuate nuove
metodologie in ricerca clinica). Ad
esempio, nel caso del glioblastoma,
grazie alla cooperazione internazionale
nell'ambito del protocollo EORTC
22981, è stato possibile ottenere
finalmente un miglioramento della
prognosi, con una nuova combinazione
di chemio-radioterapia, portando la
sopravvivenza a 2 anni dal 10 al 27%.
È stato inoltre dimostrato che la
presenza di una peculiare alterazione
genetica porta la percentuale a oltre il
40%. Quindi, pur essendo rari, i tumori
cerebrali sono una parte importantissima
della attività clinica e della ricerca
di punta nella medicina moderna.

Per un trattamento ottimale è necessaria
la collaborazione di diversi specialisti,
dal neurochirurgo, al patologo,

Studi nazionali ed internazionali

Lancet Oncology, 2012

Articles

Risk of recurrence of gastrointestinal stromal tumour after surgery: an analysis of pooled population-based cohorts

Heikki Joensuu, Aki Vechtari, Jaakko Riihimäki, Toshirou Nishida, Sonja E Steigen, Peter Brabec, Lukas Plank, Bengt Nilsson, Claudia Cirilli, Chiara Braconi, Andrea Bordoni, Magnus K Magnusson, Zdenek Linke, Jozef Sufliarsky, Massimo Federico, Jon G Jonasson, Angelo Paolo Dei Tos, Piotr Rutkowski

Summary

Background The risk of recurrence of gastrointestinal stromal tumour (GIST) after surgery needs to be estimated when considering adjuvant systemic therapy. We assessed prognostic factors of patients with operable GIST, to compare widely used risk-stratification schemes and to develop a new method for risk estimation.

Methods Population-based cohorts of patients diagnosed with operable GIST, who were not given adjuvant therapy, were identified from the literature. Data from ten series and 2560 patients were pooled. Risk of tumour recurrence was stratified using the National Institute of Health (NIH) consensus criteria, the modified consensus criteria, and the Armed Forces Institute of Pathology (AFIP) criteria. Prognostic factors were examined using proportional hazards

Published **Online**
December 7, 2011
DOI:10.1016/S1470-2045(11)70299-6

See **Online/Comment**
DOI:10.1016/S1470-2045(11)70344-8

Department of Oncology,
Helsinki University Central

Assume responsabilità e compiti nazionali ed internazionali

- Membro gruppo di lavoro EUROCARE
- Eurocourse
- Find and code
- Commissione scientifica NICER
- Executive board IARC (processo valutativo per accreditamento registri)
- European partnership against cancer

Richieste di consulenza e di dati

Registro Tumori Canton Ticino
Via in Selva, 24 - 6600 Locarno
fax: +41(0)91 816 08 29
e-mail: andrea.borsani@it.ch

FORMULARIO RICHIESTA DATI NON AGGREGATI PER PROGETTI DI STUDIO SPECIFICI
(richiesta formulabile da)

1) **Dati del Richiedente**
Cognome: ACC

Registro Tumori Canton Ticino
Via in Selva, 24 - 6600 Locarno
fax: +41(0)91 816 08 29
e-mail: andrea.borsani@it.ch

FORMULARIO RICHIESTA SCIENTIFICA

1) **Dati del Richiedente**
Cognome: Zucca
Nome: Emanuele
Istituto e indirizzo: DOSI
Recapito telefonico:
E-mail:

2) **Tipo di Richiesta Scientifica** (porre una X su una delle possibilità riportate)
 Analisi statistica
 Progetto di ricerca (studio di fattibilità, studi collaborativi)
 Formazione / aggiornamento
 Richiesta giornalistica
 Supporto scientifico
 Altro

3) **Descrizione della Richiesta**
Richiesta Hodgkin anaplastica

4) **È prevista una forma di rimborso per le prestazioni del Registro Tumori del Canton Ticino?**
 Sì
 No

Luogo e data: _____
Firma del Richiedente: _____

In caso di utilizzo delle statistiche o testi forniti, si ricorda che è auspicata la citazione della fonte: Registro Tumori Canton Ticino.

Il personale del Registro Tumori del Canton Ticino è tenuto al segreto professionale medico secondo l'articolo 321bis del Codice Penale Svizzero e la Legge Cantonale sul Registro Tumori. Pertanto, non è autorizzata alcuna forma di fornitura dati nominativi, né di singoli pazienti, né di gruppi (vedi le Basi Legali in www.it.ch/tumori/).

Registro Tumori Canton Ticino
Via in Selva, 24 - 6600 Locarno
fax: +41(0)91 816 08 29
e-mail: andrea.borsani@it.ch

FORMULARIO RICHIESTA SCIENTIFICA

1) **Dati del Richiedente**
Cognome: Manginello
Nome: Lucia
Istituto e indirizzo:
Recapito telefonico:
E-mail:

2) **Tipo di Richiesta Scientifica** (porre una X su una delle possibilità riportate)
 Analisi statistica
 Progetto di ricerca (studio di fattibilità, studi collaborativi)
 Formazione / aggiornamento
 Richiesta giornalistica
 Supporto scientifico
 Altro

3) **Descrizione della Richiesta**
Richiesta Follow up dati Centro di Senologia

4) **È prevista una forma di rimborso per le prestazioni del Registro Tumori del Canton Ticino?**
 Sì
 No

Luogo e data: 8.2.2012
Firma del Richiedente: _____

In caso di utilizzo delle statistiche o testi forniti, si ricorda che è auspicata la citazione della fonte: Registro Tumori Canton Ticino.

Il personale del Registro Tumori del Canton Ticino è tenuto al segreto professionale medico secondo l'articolo 321bis del Codice Penale Svizzero e la Legge Cantonale sul Registro Tumori. Pertanto, non è autorizzata alcuna forma di fornitura dati nominativi, né di singoli pazienti, né di gruppi (vedi le Basi Legali in www.it.ch/tumori/).

Divulgazione dati: webpage

ti Repubblica e Cantone Ticino

Organizzazione | Tematiche | Mappa | Cerca nel sito

DSS > DSP > Istituto cantonale di patologia

Registro cantonale dei tumori

www.ti.ch/tumori

► Registro cantonale dei tumori

- Presentazione
- Informazioni generali
- Metodi

- Dati generali**
- Dati specifici

- Relazioni e Posters
- Formazione
- Pubblicazioni
- Studi e progetti
- QC3 - Qualità delle Cure nella lotta Contro il Cancro

- Agenda
- Rassegna stampa
- Risorse web

In primo piano

- QC3** Progetto QC3
Qualità delle Cure nella lotta Contro il Cancro
- I tumori in Svizzera**
Situazione e sviluppi dal 1983 al 2007
- Cancer Mapping in Alpine Regions**
2001-2005

Dati generali

Dati di incidenza dei tumori

- In Ticino**
 - [Frequenza in Ticino per anno di incidenza, periodo 1996-2009.](#) (elaborati dal Registro Tumori)
 - [Trend di incidenza in Ticino e Svizzera, periodo 1996-2008.](#) (elaborati dal NICER)
 - [Alcuni approfondimenti di epidemiologia dei tumori in Ticino](#)
 - [Dati di incidenza in Ticino, periodo 1998-2002.](#) (elaborati dalla IACR. Cancer Incidence in Five Continents, vol. IX.)
 - [Dati di incidenza in Ticino, periodo 1998-2002, per istotipo, uomini.](#) (elaborati dalla IACR. Cancer Incidence in Five Continents, vol. IX.)
 - [Dati di incidenza in Ticino, periodo 1998-2002, per istotipo, donne.](#) (elaborati dalla IACR. Cancer Incidence in Five Continents, vol. IX.)
- In Svizzera**
 - [Dati elaborati dal National Institute for Cancer Epidemiology and Registration \(NICER\)](#)
- Nel Mondo**
 - [Cancer Incidence in Five Continents, volume IX. Dati elaborati dalla International Association of Cancer Registries \(IACR\)](#)

Dati di mortalità dei tumori

- In Ticino**
 - [Trend di mortalità in Ticino e Svizzera, periodo 1984-2008.](#) (elaborati dal NICER)
- In Svizzera**
 - [Dati elaborati dal National Institute for Cancer Epidemiology and Registration \(NICER\)](#)

Dati di Sopravvivenza dei pazienti ticinesi per tipologia tumorale

AC: adenocarcinoma; LLC: large cell carcinoma; SmCC: small cell carcinoma;
SqCC: squamous cell carcinoma

QC3 project

Tab. 3 – SELEZIONE DEGLI IQ RIGUARDANTI I CASI DI TUMORE COLO-RETTALE INCIDENTI NEL 2011

INDICATORE (IQ)	DENOMINATORE	SI	NO	NR	P	CCS	R	G	RCT	M	PBS
Proportion of patients evaluated by preoperative colonoscopy	Patients with colorectal cancer undergoing surgery (n=200)	86%	8.5%	5.5%		*	*	**			
Proportion of patients with preoperative staging according to the AJCC TNM 7 th ed	Patients with colorectal cancer undergoing surgery (n=200)	63%	37%	0%				*			
Proportion of patients undergoing rectal-sigmoidoscopy /colonoscopy	Patients with rectal cancer (n=76)	89.5%	4%	6.5%	*	*	**			*	
Proportion of patients with a visit, particularly during the last year of life											
Proportion of patients including the last visit											
Proportion of patients including the last visit and the last hospitalization											
Proportion of patients including the last visit and the last hospitalization and the last surgery											
Proportion of patients with a number of visits greater than 1											
Proportion of patients (every T, N1-2, M0) with anastomosis											
Proportion of patients with systemic therapy (KRAS, BRAF)											
Proportion of patients with hepatic metastases undergoing immediate/synchronous metastasectomy	singular pulmonary metastases (n=30)										
Proportion of patients with protective stoma before neo-adjuvant RT-CT	Patients with rectal cancer undergoing neo-adjuvant radio-chemotherapy (n=20)	10%	80%	10%		**					
Proportion of patients with locally advanced tumours undergoing neo-adjuvant RT±CT	Patients with locally advanced rectal cancer (n=24)	70.8%	25%	4.2%		****	**	**			

Ongoing work: Survival
 QC3: indicatori di qualità delle cure oncologiche in Ticino 2011-2013

Screening mammografico

British Journal of Cancer (2009), 1–7
© 2009 Cancer Research UK. All rights reserved 0007–0920/09 \$32.00
www.bjcancer.com

Full Pa
Asse
clinic

A B

Ticino
ISPM
Switze

BACK
the d
scree
METH
Switz
diane
RESUL
increa
decre
tum
Chan
CONC

Ongoing work: Survival Screening mammografico

are still less favourable than those achieved by organised screening programmes.
British Journal of Cancer advance online publication, 27 October 2009; doi:10.1038/sj.bjc.6605378 www.bjcancer.com
© 2009 Cancer Research UK